


NAVAL OCEANOGRAPHIC OFFICE

SURFACE CURRE

SOUTHEAST INDIAN OCEAN IN THE GREAT AUSTRALIAN B



OCTOBER 1977 REPRINTED 1980



APPROVED FOR PUBLIC SELECTE DISTRIBUTION ON


DEPARTMENT OF THE NAVY NSTL STATION, MISSISSIPPI 395 AL OCEANOGRAPHIC OFFICE SPECIAL PUBLICATION 1404—IN 8

URFACE CURRENTS

(2)

HEAST INDIAN OCEAN INCLUDING HE GREAT AUSTRALIAN BIGHT

LEVELI



OCTOBER 1977
REPRINTED 1980



Serent D

SURFACE CURRENTS.

SOUTHEAST INDIAN OCEAN INCLUDING THE GREAT AUSTRALIAN BIGHT.



9 = 1, my

REPRINTED 1980

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

NAVAL OCEANOGRAPHIC OFFICE NSTL STATION, MISSISSIPPI 39522

URFACE CURRENTS.

HEAST INDIAN OCEAN INCLUDING HE GREAT AUSTRALIAN BIGHT



9 = 1, 11/1

435

∫; ØCT**≎≎€₹≈**₹977

REPRINTED 1980

POVED FOR PUBLIC RELEASE, DISTRIBUTION UNLIMITED

NAVAL OCEANOGRAPHIC OFFICE NSTL STATION, MISSISSIPPI 39522

ACKNOWLEDGMENTS

Messrs. Raymond J. Beauchesne \star and William E. Boisvert made major contributions to this atlas.

^{*}Mr. Beauchesne presently is employed by the Bureau of Naval Personnel.

FOREWORD

THIS ATLAS, ONE IN A SERIES OF 43 REGIONAL SURFACE CURRENT ATLASES, IS PRODUCED TO FULFILL A NEED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC AND INDUSTRIAL COMMUNITIES FOR THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA. THESE ATLASES ADD TO THE WEALTH OF NAUTICAL INFORMATION UPON WHICH OPERATIONAL PLANNING, NAVIGATIONAL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID PRODUCTION AND WIDE DISSEMINATION OF THIS ATLAS ARE MADE POSSIBLE BY THE LATEST COMPUTER TECHNIQUES.

THE CONSTANT IMPROVEMENT IN THE QUALITY OF SURFACE CURRENT DATA RECEIVED OVER THE YEARS IS MADE POSSIBLE LARGELY BY THE MORE THOROUGH REPORTS OF VOLUNTARY OBSERVERS IN RECENT YEARS. THE DEFENSE MAPPING AGENCY, THE OCEANOGRAPHIC OFFICE, AND THE USER OF THE ATLASES RELY ON THE PERSONAL OBSERVATIONS OF THE MAN WHO HAS "BEEN THERE." MARINERS, IN REPORTING THEIR OBSERVATIONS, RENDER A SERVICE NOT ONLY TO THEMSELVES BUT ALSO TO ALL "WHO GO DOWN TO THE SEA IN SHIPS." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONIC NAVIGATION AIDS, AND 300,000-TON SHIPS, UP-TO-DATE, RAPIDLY DISSEMINATED ENVIRONMENTAL AND NAVIGATIONAL INFORMATION HAS BECOME INCREASINGLY IMPORTANT.

JOHN R. McDOWNELL Captain, U.S. Navy

Commander

Acce MIII DTI Una-Just

By . Dist

Dist

A

FOREWORD

N A SERIES OF 43 REGIONAL SURFACE CURRENT ATLASES. ED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC AND THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA. ALTH OF NAUTICAL INFORMATION UPON WHICH OPERA-AL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID NATION OF THIS ATLAS ARE MADE POSSIBLE BY THE

DVEMENT IN THE QUALITY OF SURFACE CURRENT DATA MADE POSSIBLE LARGELY BY THE MORE THOROUGH REPORTS ECENT YEARS. THE DEFENSE MAPPING AGENCY, THE HE USER OF THE ATLASES RELY ON THE PERSONAL OB-AS "BEEN THERE." MARINERS, IN REPORTING THEIR ICE NOT ONLY TO THEMSELVES BUT ALSO TO ALL "WHO ." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONIC }-TON CHIPS, UP-TO-DATE, RAPIDLY DISSEMINATED TAL INFORMATION HAS BECOME INCREASINGLY IMPORTANT.

> JOHN R. McDORWELL Captain, U.S. Navy

Commander


Acce	acton For
DTIC	
Unan Just	nounced ification
Ву	
	ribution/
Ava	llability Codes
	Avail and/or
Dist	Special
Δ	

SURFACE CURRENT ATLASES

THIS GERIES FORMPUTERIZED ATTRIES REPUBLIES THE TO BE RECRAPHT FOR A STATES FOR THE CURRENTS CHOP 366, 368, 369, 700 AHT HAFRE MANUALLY OMPTION FROM DATA OBTAINED DURING THE PERIOD 1993 - 1994. THISS NEW ATTRASES ONE RM TO THE STANDARD NATY OCEAN AREA AND REGION INDEX CIMITS SHOWN BELOW: HOLD, NOT STANDARD TO VERY NORTH PACIFIC REGION TO EAST OF THE PHILIPPINES.

AND SHOULD BE AND A STANDARD BY A STANDARD B

ROUGHT IMPROVEMENTS IN THE DATA FOR ASSURE THE INCOMING OF THE DATASE, HIGH CHALLTY SURFACE DEBRENT DATA ADAILABLE. THE FILE NOW CONTAINS MORITHAN (1.00, 100, 100, 18SERVATIONS AND A GENERAL PRATE OF THE FILE WILL BE MADE




SURFACE CURRENT ATLASES

FY RUGBAPHI FEIGE WERE MANEALLY THESE NEW ATLASES IMITS SHOWN BELOW: THE THE PHILIPPINES.

DUNCE THE DATE TO A TENTAL OF A NUMBER OF THE SECOND SECON

NO AMEN'NTS OF NEW LAIR WARRANT, MODILING BOOKERS OF SIMPLANTING



Introduct ion

The Surface Gutrent Data File, from which these stlames are derived, consists orlearly of over four million ship ser and drift observations. These data were collected by the Netherlands, Japan, Britain, France, and the inited States. The file is supplemented by several thousand Geomagnetic Fleetrokinetograph (GEK) observations, mostly impaness. The file spans the period from the early 1850's to the present. The earliest observations were collected by the Netherlands and Great Britain; those of the 1960's through the present are primarily inited States data.


General Quality

The quality of this data five is considered high for this type of derived value. The data have been carefully screened for duplication; observations taken under adverse conditions (i.e. high winds and waves, time between observations greater than 1, nourse have been eliminated when warranted, consideration was given to the reliability of the observer; doubtill shipboard commutations of set and frift were edited, and observations with erroneous locations (mostly observations on land, have been eliminated. The accepted data are considered most iseful when used collectively as in summaries where a number of observations show trends.

General Observation Technique

The set (direction) and srift (speed) are computed by the navigator from the difference between the dead reckoning (DR) position and the position determined by any type of navigational fix. The iritrian be determined along any straight line traik and includes all factors which cause changes in the DR position. When a fix is obtained, the urrent set (direction) is FROM the DR position. To the fix, the drift (speed) is equal to the distance in nautical miles between the DR and the fix, divided by the number of hours since the last fix. For successive observations, the TO POSITION of one observation becomes the FROM POSITION of the pext observation.


Because the influence of current may vary along a ship's track, the MEAN POSIT ON of the track is assigned as the geographic location of the current observation. An example of a current computation is shown in the figure below.




EXAMPLE OF A SURFACE CURRENT (SHIP'S DRIFT) OBSERVATION

Data Presentation


The following legend shows two types of surface current presentations by 1° quadrangle, type 1 with 12 or more observations and type 2 with fewer than 12 observations. Where there are 11 or fewer observations within a 1° quadrangle, the total number of observations is shown within the 90° quadrant containing the observations.



It chepe are I se wire chart a no sector lessitants as the wi



(D) Persister Carrent 50 percent in more or a frame of 401 chaecarions (a) within a 40° section (b) of the Bepoint compass.



(4) <u>Bizonal Flow</u> - Practically all observation are concentrated in apposite pairs of a sectors, and one pair contains at least 80 percent as many observations as the oppair. This generally indicates variabilithat occurs in zones of entrainment between the containment of the co



of these ase of impose inservarious on a figurationage in elegenous encountries of some objection of the contributions of the contributions of the contributions as to come

majerico di Properti di Proper

Silver Si

Control Section (Control Section (Contro

i **A** i ir examb e



 Pensianer urben 50 gehrent tragere 30 sul harrances fati wiretnia 20 gehran nitse 8-bught nempasa.




7. Prevailing Gurrent 10 percent or more of all chemications fail within two adjacent 20° sectors.



a Primary Turnenn with Gelondary lyne cum a Primary Largent - 10 percent or ever of all Inservations fall within three adjacent 49° sectors.

b Secondary Offertion - 25 percent of more of all observations fall within a 45° sector, and the two resultant vector directions are separated by more than 36 of arc.




sty Bizonal Flow - Practically all observations are subpentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).




(5) Variable Current - The 45° sector with most observations has less than 25 percent of all observations; direction is indeterminate.


adrangie Where Theorya


A CONTRACT OF THE CONTRACT OF





---)


























DISTRIBUTION LIST

NAVY

CINCPACELT (02M) COMTHIRDFLT COMSEVENTHELT COMSUBPAC COMNAVAIRPAC COMPATWINGSPAC PATWINGSPAC DET ADAK PATWING 1 COMNAVSURFPAC DI RNAVOCEANMET FLENUMEACEN FLEWEACEN GUAM FLEWEACEN PEARI. NAVWEASERVFAC SAN DIEGO NAVWEASERVFAC YOKOSUKA NWSD ASHEVILLE NWSED ADAK NWSED AGANA NWSED ATSUGI NWSED KADENA NWSED MISAWA

OTHER GOVT.

NOAA/NODC NOAA/NCC

PRIVATE & UNIVERSITIES

FLORIDA ST. UNIV.
LOUISIANA ST. UNIV.
MASS. INST. OF TECH
ORE. ST. UNIV.
TEXAS A&M UNIV.
UNIV. OF MIAMI
UNIV. OF R.I.
UNIV. OF WASH.
SCRIPPS INST OF OCEANOGRAPHY
WOODS HOLE OCEANOGRAPHIC INST.

FOREIGN

HYDROGRAPHER/R.A.W.
DEPT. TRANSPORTATION/AUSTRALIA

REPORT DOCUMENTATION PAGE	REAT INSTENDED TONG BEFORE TMELATING A RM
113101.14310	MAN WOLL - BET - BUT S TOTA 13 V. WEES
AD-A093	
A CONTROL OF THE PROPERTY OF T	4 TYPE OF BEDOM THE PERSON TO FAIR
- 11 (94) (1 1 22%) - 1 - 11 (24) - 18 (24) - 21 (25) (27)	75.11
	s Remesam us res movement ungen
1776,	के गण्यानकशान एक प्रकार स्थानिक के व
- NA Francis (near ver light lift) re - Na Francis (no 1990)	Į.
A Section of the Control of the Cont	i
FORDE MING DROAN EAT ON NAME AND ADDRESS	'ଓ ୨୭୦୯୯ଟ ଅଟେ ଅନ୍ତର୍ଗିତ " ୨୬୯୯ ଜଳରେ ବ୍ୟତ୍ତିକ ଓଡ଼ି ବ୍ୟକ୍ତିକ୍ତି
Naval indonaracis Office	
TONTED IN NO SERVICE NAME AND ADDRESS	1 eseger sars
	13 M. MRER OF PAGES
	ात के कि कि एन में बहुर हैं। वि
I MON "SENS ASEN" VAME & ACCRESSITE STORES From Control ing	Office 15 SECURITY CLASS of this report.
	, <u> </u>
	191 191 191 191 191 191 191 191 191 191
	SE DESLASSIFICATION DOWN SRADING
Approved for public release; distribution on	nlim(tet-
: DISTRIBUTION STATEMENT of the abstract entered in Block 20. If di	
**: DISTRIBUTION STATEMENT of the abetract entered in Block 20. If di	Tarant (from Report,
4 S. PPLEMENTARY NOTES	
4 STALTEMEN MAY 40.52	1
	1
19 KEY WORDS (Continue on reverse side it necessary and identify by bloc	k number)
	i
Surface Currents Southeast Indian Ocean	
Great Australian Bight	í
The constitution of the co	1
20 ABSTRACT (Continue on reverse olds if necessary and identify by black	
This atlas, and the series of which it i	s a part, is computer generated
and automatically plotted. It makes availab	
surface current data collected and will be u of data are added to the data file. This an	
on a vast quantity of data as compared to th	
editions printed in the mid-thirties.	

The second secon

DO 170 1473 EDITION OF 1 NOV 84 IS OBSOLETE S/N 0102-014-8601

SECURITY CLASSIFICATION OF THIS PAGE (When Date Secured)

SECURITY SUBSEIFICATION "

	Commence and the second of the
PEAG INSTRUCTIONS	h to the agree of a transplace of the Park Park Data Entered
READ INSTRUCTIONS REFORE COMPLETING FORM THEFATS CATALOG NUMBER	
	2 Cont.
PT SE MEMORY & PEMOG COVERED	The surface current information is based majory in solo brift, which
<u> </u>	is the difference between the dead reckening position and the position determined by any type of navigational fix. This difference describes
FFSHMINS THE REPORT NUMBER	the direction and speed of the current.
FRACT OR TRANT HUMBERIA	*
}	
IGGRAM F. EMENT PROJECT TASK	
TEA & BOTH JAIT HUMBERS	
PORT 34"5	
WATE OF PAGES	
SURITY CLASS (of this report)	
FOL ASSIF CA TON DOWNGRADING	
HEDULE	
\	
	}
}	}
j	
1	1
	\
computer generated	
he most recent ver sufficient amounts	1
atlases are based anually-compiled	
and the completed	
1	1

IN ST THIS PARE PAREN BUT BARRES

SECURITY CLASSFICATION OF THIS PASS/When Date British

DATE ILMED